

CLAIMS

We claim:

1. A combination of antichlamydial agents comprising at least two agents, each of which is effective against a different phase of chlamydial life cycle.
2. The combination according to Claim 1 wherein the agents are selected from the group consisting of:
 - a) agents effective against cryptic phase of chlamydial life cycle;
 - b) agents effective against elementary body phase of chlamydial life cycle; and
 - c) agents effective against replicating phase of chlamydial life cycle.
3. The combination according to Claim 1 wherein the agents are assembled as an admixture.
4. The combination according to Claim 1 wherein the agents are copackaged individually.
5. The combination according to Claim 1 wherein the agents are instructionally assembled.
6. The composition according to Claim 1 further comprising antiinflammatory agent, immunosuppressive agent, Vitamin C or combinations thereof.
7. A combination of antichlamydial agents for use in managing chlamydial infection or prophylaxis thereof; wherein the combination comprises at least two agents, each of which is effective against a different phase of chlamydial life cycle.

067097-1080

8. method of treating a chlamydial infection or disease associated therewith comprising treating the patient to reduce the acellular load of infectious EBs.
9. A method of treating a chlamydial infection or disease associated therewith comprising treating the patient with one or more antibiotics for not less than 45 days.
10. A pharmaceutical composition comprising at least two agents, each of which is effective against a different phase of chlamydial life cycle.
11. The pharmaceutical composition according to Claim 10 wherein the agents are selected from the group consisting of:
- a) agents effective against cryptic phase of chlamydial life cycle;
 - b) agents effective against elementary body phase of chlamydial life cycle; and
 - c) agents effective against replicating phase of chlamydial life cycle.
12. The pharmaceutical composition according to Claim 10 wherein the agents are formulated together in a physiologically acceptable vehicle.
13. A pharmaceutical composition for use in managing chlamydial infection or prophylaxis thereof; wherein the composition comprises at least two agents, each of which is effective against a different phase of chlamydial life cycle.

00307-10300

14. The pharmaceutical composition according to Claim 13 further comprising antiinflammatory agent, immunosuppressive agent, Vitamin C or combinations thereof.
- 5 15. A method for eliciting a protective immune response to *Chlamydia* infection in an animal or human by administering to an infected animal or human at least two agents, each of which is effective against a different phase of chlamydial life cycle until the human or animal no longer tests positive for *Chlamydia*.
- 10 16. A diagnostic kit or pack comprising materials adequate to perform one or more of the following:
- 15 a) antibody assays against recombinant MOMP;
- b) antibody assays against specific antigenic peptides described in Figures 3, 4 and 5;
- c) antigen capture assays directed against MOMP or the peptides described in Figures 3, 4 and 5;
- 20 d) DNA amplification assays for chlamydial genes; and
- e) Western blot used as a confirmatory.
17. The kit or pack of Claim 16, wherein the antibody assays are ELISAs.
18. A pharmaceutical pack of therapeutic agents for management of *Chlamydia* infection, comprising at least two agents, each of which is effective against a different phase of chlamydial life cycle.
- 25 19. The pharmaceutical pack according to Claim 18 wherein the agents are selected from the group consisting of:

0970924110300

- 5
- a) agents effective against cryptic phase of the chlamydial life cycle;
 - b) agents effective against elementary body phase of the chlamydial life cycle; and
 - c) agents effective against replicating phase of the chlamydial life cycle.

20. The pharmaceutical pack according to Claim 18 further comprising antiinflammatory agent, immunosuppressive agent, Vitamin C, or combinations thereof.

10 21. The pharmaceutical pack according to Claim 18 wherein the pack is a single unit dose.

22. The pharmaceutical pack according to Claim 18 wherein the agents are contained within the pack separately and/or as an admixture.

15 23. The pharmaceutical pack according to Claim 18 wherein the pack is a single blister pack or a single vial.

24. The pharmaceutical pack according to Claim 18 wherein the pack comprises a plurality of unit dosages.

20 25. The pharmaceutical pack according to Claim 18 wherein the pack is a plurality of single blister packs or a plurality of single vials.

26. A method for treating biological material infected with *Chlamydia*, comprising contacting the biological material with at least two agents, each of which is effective against a different phase of chlamydial life cycle.

25

008071026060

-86-

- 00307 " 110300
27. The method according to Claim 26 wherein the agents are selected from the group consisting of:
- a) agents effective against cryptic phase of chlamydial life cycle;
 - 5 b) agents effective against elementary body phase of chlamydial life cycle; and
 - c) agents effective against replicating phase of chlamydial life cycle.
28. The method according to Claim 26 wherein the agent effective against the elementary body phase is a disulfide reducing agent.
29. The method according to Claim 29, wherein the agent effective against the cryptic phase is a nitroaromatic compound.
- 15 30. The method according to Claim 26, wherein the nitroaromatic compound is selected from the group consisting of nitroimidazoles, nitrofurans, analogs, derivative and combinations thereof.
31. A method for managing chlamydial infection in an individual in need thereof, comprising administering at least two agents, each of which is effective against a different phase of chlamydial life cycle.
- 20 32. The method according to Claim 31 wherein the agents are selected from the group consisting of:
- 25 a) agents effective against cryptic phase of chlamydial life cycle;
 - b) agents effective against elementary body phase of chlamydial life cycle; and

1

- c) agents effective against replicating phase of chlamydial life cycle.
33. A method for diagnosing disease associated with *Chlamydia* infection comprising diagnosing the *Chlamydia* infection in an individual, wherein the disease is an autoimmune disease, an inflammatory diseases or a disease that occurs in immunocompromised individuals.
34. A method for treating disease associated with *Chlamydia* infection comprising treating the *Chlamydia* infection in an individual in need thereof, comprising administering a combination of antichlamydial agents that are effective against at least two phases of the chlamydial life cycle. wherein the disease is an autoimmune disease, an inflammatory diseases and disease that occurs in immunocompromised individuals.
35. The method according to Claim 34 wherein the agents are selected from at least two different agents from at least two of the following groups:
- a) agents effective against cryptic phase of chlamydial life cycle;
 - b) agents effective against elementary body phase of chlamydial life cycle; and
 - c) agents effective against replicating phase of chlamydial life cycle.
36. The method according to Claim 35 wherein the infection is a *Chlamydia pneumoniae* infection.

37. An assay for identifying an agent which is capable of inhibiting chlamydial infection, comprising the steps of:

- a) preparing tissue culture cells infected with *Chlamydia* in the absence of cycloheximide;
- b) allowing the *Chlamydia* to replicate;
- c) adding a test agent;
- d) isolating chlamydial nucleic acid from the cells;
- e) amplifying the chlamydial nucleic acid by a nucleic acid amplification technique; and
- f) evaluating the presence or absence of amplified chlamydial nucleic acid;

wherein the absence of amplified chlamydial nucleic acids is indicative that the agent is capable of inhibiting chlamydial infection.

38. The assay of Claim 37 wherein the amplification technique is PCR.

39. A method of identifying cells containing cryptic form of *Chlamydia* comprising the steps of:

- a) treating cultured cells, thought to be infected with chlamydia, with a disulfide reducing agent;
- b) subjecting cultured cells to protease digestion;
- c) exposing cells to appropriate polymerase, dNTPs and primers for DNA amplification of a chlamydial protein;
- d) exposing the cells to a reporter molecule enzyme;
- e) exposing the cells to an appropriate substrate for the reporter enzyme; and
- f) determining the presence of cryptic form of *Chlamydia* by visualizing the amplified DNA encoding a chlamydial protein.

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

- DELETED COPY**

-90-

- g) exposing the cells to an appropriate substrate for the reporter enzyme;
- h) determining the presence of cryptic form of *Chlamydia* by visualizing the amplified DNA encoding a chlamydial protein;

wherein the absence of amplified chlamydial nucleic acids is indicative that the agent is capable of inhibiting chlamydial infection.

44. An assay according to Claim 43, wherein the appropriate primers of step a) are CHLMOMPDB2 and CHLMOMPCB2.
45. A method for activating macrophages or monocytes in which their infection fighting ability has been compromised by a *Chlamydia* infection, comprising treating the *Chlamydia* infection by contacting the infected macrophages or monocytes with an antichlamydial agent.
46. A method of treating biological material infected with *Chlamydia*, comprising contacting the biological material with at least one agent selected from the following groups:
- a) agents effective against the cryptic phase;
 - b) agents effective against elementary body phase;
 - c) ethambutol and anules and derivatives thereof and isonicatinic acid cogeners;
 - d) agents first identified as inhibiting chlamydial infection by the method of Claim 36.
47. A method of detecting chlamydial elementary bodies in a sample comprising contacting the sample with a disulfide reducing agent before using a DNA

a
amplification technique to detect chlamydial DNA in the sample.

48. A method of detecting chlamydial elementary bodies in a sample comprising performing on the sample an antigen capture assay directed against MOMP or the peptides described in Figures 3, 4 and 5.

49. The method of Claim 48, wherein the sample is contacted with a disulfide reducing agent before the assay is performed.

10 50. A peptide of Figure 3, 4 and 5.

51. Antichlamydia agent identified by the assay of Claim 37.

52. Antichlamydia agent identified by the assay of Claim 43.

15 53. The method of Claim 46 wherein the agent effective against the cryptic phase is a nitroaromatic compound.

54. The method of Claim 46 wherein the agent effective against the elementary body phase is a disulfide reducing agent.

20
25 55. A method of determining the status of a patient or monitoring and/or modifying the course of therapy for chlamydia infection comprising the results of one or more assays made contemporaneously or sequentially, wherein the assays are selected from the group consisting of:

-92-

- 5 a) antibody assays against recombinant MOMP;
b) antibody assays against specific antigenic
peptides described in Figures 3, 4 and 5;
c) antigen capture assays directed against MOMP or
the peptides described in Figures 3, 4 and 5;
d) DNA amplification assays for chlamydial genes;
and
e) Western blot used as a confirmatory.
- 10 56. The method of Claim 55, wherein the antibody assays
are ELISAs.
57. The method of Claim 55 wherein DNA amplification
assay is PCR.
58. The method of Claim 55 wherein DNA amplification
assay amplifies the MOMP gene.
- 15 59. The method of Claim 55 wherein more than one assay is
used.
60. The method of Claim 55 wherein specific antibody
measures are obtained against IgA, IgE, IgM and/or
IgG.
- 20 61. An assay for identifying an agent which is capable of
inhibiting chlamydial infection in a patient or an
animal, comprising the steps of:
- 25 a) obtaining a biological material sample from a
patient from which the infectious organism is
extracted and applied to target cells for
culturing;
b) allowing the *Chlamydia* to replicate;
c) adding one or more agents;

008077-110800

5

10

10

10

15

15

15

□

□

10

a) obtaining serum or whole blood from an animal or human that has undergone therapy for eliminating

070901-1080

Chlamydia infection; wherein the therapy comprises administering to the animal or human at least two agents effective against a different phase of the chlamydial life cycle;

- 5 b) introducing the serum or whole blood into an *Chlamydia*-free organism; and
- c) determining bacterial response to chlamydial challenge in the *Chlamydia*-free organism.

add
G3

008011-110300